

Non-zero Fermion Modes Contribution to the Instanton-Induced Electroweak Processes at High Energies

R.G.Shulyakovsky

*B.I. Stepanov Institute of Physics,
National Academy of Sciences of Belarus
F. Skorina avenue 68,
220072 Minsk, Belarus
E-mails: shul@dragon.bas-net.by*

We consider instanton-induced high energy processes in an electroweak-type model. Multi-fermion final states are taken into account in Gauss approximation. Unlike previous results we calculate nonzero fermion modes contribution to cross-section by straightforward procedure. This contribution can be important at energies of sphaleron mass order.

Key words: instantons, non-zero fermion modes, baryon number violation

PACS numbers: 11.15.Tk

1 Introduction

As is well known [1] instanton-induced processes in electroweak theory can violate baryon number and concern the problem of matter and antimatter asymmetry of the Universe [2]. At the beginning of preceding decade it was shown [3, 4, 5] that cross-section of these processes exponentially increase with energy (see for review [6])

$$\sigma^{inst}(\varepsilon) \sim e^{\frac{16\pi^2}{g^2}F(\varepsilon)}, \quad F(\varepsilon) = -1 + \frac{9}{8}\varepsilon^{\frac{4}{3}} - \frac{9}{16}\varepsilon^2 + \dots, \quad \varepsilon = \frac{\sqrt{s}}{E_{sp}}, \quad (1)$$

where $\sqrt{s}$ is c.m. energy of the process, $E_{sp} \approx \frac{4\sqrt{6}\pi m_w}{g^2} \approx 16 \text{ TeV}$ - energy of sphaleron [7]. The first term in "Holy Grail" function $F(\varepsilon)$ is 't Hooft suppression factor [1], the second corresponds to the multiple W - and Higgs bosons production in Gauss approximation [4, 5], the third is a contribution of tree boson graphs [8, 9]. Boson loops [10] and fermion contribution [11] are accumulated in pre-exponent factor. One can naively suppose that $F(\varepsilon) \rightarrow 0$ if $\sqrt{s} \rightarrow E_{sp}$ and cross-section (1) can reach observable value ¹. In this case pre-exponent constant in (1) can be important.

¹Exact behavior of "Holy Grail" is unknown at $\varepsilon \geq 1$ and a question about possibility of observation of anomalous baryon number violation is still opened.

Contribution of W- and Higgs bosons diagrams to electroweak instanton processes was calculated in detail (see f.e. [3]-[6],[8]-[10]) whereas graphs with arbitrary number of fermion legs were rarely taken into account. In the pioneer paper of O.Espinosa [11] author found multi-fermion contribution by means of coherent-state formalism. In this paper we are checking the results of Ref. [11] by straightforward procedure in Gauss approximation and also finding the distribution on fermion multiplicity.

2 Multi-fermion Instanton-Induced Green Functions

Let us consider electroweak-type model in the limit of vanishing Weinberg angle with the following Euclidean action (we will present most formulas in Euclidean space; analytical continuation to the Minkovsky space will be performed in the final step)

$$S[A, \psi, \bar{\psi}] = S_{YM} + S_f + S_H, \quad S_{YM} = \frac{1}{2} \int dx \text{Tr} (F_{\mu\nu} F_{\mu\nu}), \quad (2)$$

$$S_f = - \int dx \sum_{i=1}^{n_f} \bar{\psi} i \not{D} \psi, \quad S_H = \int dx \left((D_\mu \phi)^\dagger D_\mu \phi + \lambda \left(\phi^\dagger \phi - \frac{v^2}{2} \right)^2 \right),$$

$$\not{D} = (\partial_\mu - ig A_\mu) \sigma_\mu, \quad F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu - ig [A_\mu, A_\nu], \quad A_\mu = A_\mu^a \frac{\sigma_a}{2},$$

where n_f is a number of massless Weyl fermion doublets, $n_f = 12$ for three generations.

Let us consider for definiteness instanton-induced fermion collisions with arbitrary number of fermions, W- and Higgs bosons in final states

$$f + f \rightarrow (2m - 2)f + n_w W + n_h H. \quad (3)$$

General form of Euclidean Green function for these processes ² can be given by functional integral on gauge fields with one unit of topological charge

$$\int D A D \phi D \psi D \bar{\psi} e^{-S} \prod_{i=1}^m \psi(x_i) \prod_{j=m+1}^{2m} \bar{\psi}(x_j) \prod_{k=1}^{n_w} A_{\mu_k}(y_k) \prod_{l=1}^{n_h} \phi(z_l), \quad (4)$$

where group and spinor indexes are omitted for simplicity.

As is well known in Gauss approximation integration in (4) on boson and fermion fields can be performed independently

$$\int d^4 z \int_0^\infty \frac{d\rho}{\rho^5} \int dU \mu(\rho) e^{-\frac{8\pi^2}{g^2} - \pi^2 \rho^2 v^2} \prod_{k=1}^{n_w} A_{\mu_k}^{inst}(y_k - z) \prod_{l=1}^{n_h} \phi^{inst}(z_l - z) \times$$

$$\times \int D \psi D \bar{\psi} e^{\int dx \bar{\psi} i \not{D} [A^{inst}] \psi} \prod_{i=1}^m \psi(x_i - z) \prod_{j=m+1}^{2m} \bar{\psi}(x_j - z), \quad (5)$$

²Only one sort of fermions is taken into account. Generalization into $n_f = 12$ is trivial.

where z_μ , ρ , U and $\mu(\rho)$ are instanton center, size, orientation and density correspondingly [1, 12], A^{inst} and ϕ^{inst} are instanton solutions [13].

Since boson part of (5) was investigated quite well let us concentrate on fermion contribution

$$\langle \psi(x_1) \dots \bar{\psi}(x_{2m}) \rangle^{inst} \equiv \int D\psi D\bar{\psi} e^{\int dx \bar{\psi} i \not{D} [A^{inst}] \psi} \prod_{i=1}^m \psi(x_i) \prod_{j=m+1}^{2m} \bar{\psi}(x_j), \quad (6)$$

which can be considered as a fermion Green function in *external* instanton field.

The simplest non-vanishing Green function for only one fermion doublet reads [1]

$$\begin{aligned} \langle \psi(x_1) \bar{\psi}(x_2) \rangle^{inst} &= \prod_n \int dc_n d\bar{c}_n e^{\lambda_n c_n \bar{c}_n} \sum_s c_s \psi_s(x_1) \sum_r \bar{c}_r \bar{\psi}_r(x_2) = \\ &= \det'(\rho) \psi_0(x_1) \bar{\psi}_0(x_2), \quad \det'(\rho) = \prod_{n \neq 0} \lambda_n, \quad i \not{D} [A^{inst}] \psi_n(x) = \lambda_n \psi_n(x), \end{aligned} \quad (7)$$

where Grassman algebra is used. Explicit expressions for zero mode $\psi_0(x)$ (eigen function belonging to $\lambda_0 = 0$) and $\det'(\rho)$ are known [1].

Let us calculate 4-fermion Green function in the instanton field

$$\begin{aligned} \langle \psi(x_1) \bar{\psi}(x_2) \psi(x_3) \bar{\psi}(x_4) \rangle^{inst} &= \prod_n \int dc_n d\bar{c}_n e^{\lambda_n c_n \bar{c}_n} \psi(x_1) \bar{\psi}(x_2) \psi(x_3) \bar{\psi}(x_4) = \\ &= \int dc_0 d\bar{c}_0 c_0 \psi_0(x_1) \bar{c}_0 \bar{\psi}_0(x_2) \prod_{n \neq 0} \int dc_n d\bar{c}_n (1 + \lambda_n c_n \bar{c}_n) \psi(x_3) \bar{\psi}(x_4), \\ \prod_{n \neq 0} \int dc_n d\bar{c}_n (1 + \lambda_n c_n \bar{c}_n) \psi(x_3) \bar{\psi}(x_4) &= \sum_{k \neq 0} \left(\int dc_k d\bar{c}_k c_k \psi_k(x_3) \bar{c}_k \bar{\psi}_k(x_4) \times \right. \\ &\times \left. \prod_{n \neq 0, k} \int dc_n d\bar{c}_n \lambda_n c_n \bar{c}_n \right) = \sum_{k \neq 0} \left(\psi_k(x_3) \bar{\psi}_k(x_4) \prod_{n \neq 0, k} \lambda_n \right) = S^{nz}(x_3, x_4) \det'(\rho), \end{aligned} \quad (8)$$

where non-zero fermion modes propagator $S^{nz}(x, y)$ can be calculated analytically [14, 15]:

$$\begin{aligned} S^{nz}(x, y) &\equiv \sum_{j \neq 0} \frac{\psi_j(x) \bar{\psi}_j(y)}{\lambda_j} = \frac{1}{\sqrt{1 + \rho^2/x^2}} \frac{1}{\sqrt{1 + \rho^2/y^2}} \left[\frac{(x-y)_\mu \sigma_\mu}{2\pi^2 (x-y)^4} \times \right. \\ &\times \left(1 + \rho^2 \frac{x_\nu \sigma_\nu y_\kappa \bar{\sigma}_\kappa}{x^2 y^2} \right) - \frac{\rho^2}{4\pi^2 (x-y)^2 x^2 y^2} \times \\ &\times \left(\bar{\sigma}_\mu \frac{x_\nu \sigma_\nu \bar{\sigma}_\mu (x-y)_\lambda \sigma_\lambda y_\omega \bar{\sigma}_\omega}{\rho^2 + x^2} + \sigma_\mu \frac{x_\nu \sigma_\nu (x-y)_\lambda \bar{\sigma}_\lambda \sigma_\mu y_\omega \bar{\sigma}_\omega}{\rho^2 + y^2} \right) \Big], \end{aligned} \quad (9)$$

we use standard notations in Euclidean space $\sigma_\mu = (-i\sigma_a, I)$, $\bar{\sigma}_\mu = (i\sigma_a, I)$. The propagator can be generalized to arbitrary instanton position and color orientation.

Finally, for the 4-fermion Green function we obtain

$$\langle \psi(x_1) \bar{\psi}(x_2) \psi(x_3) \bar{\psi}(x_4) \rangle^{inst} = \psi_0(x_1) \bar{\psi}_0(x_2) S^{nz}(x_3, x_4) \det'(\rho). \quad (10)$$

Higher Green functions are calculated similarly:

$$\begin{aligned} & \langle \psi(x_1)\bar{\psi}(x_2)\dots\psi(x_{2m-1})\bar{\psi}(x_{2m}) \rangle^{inst=} \\ & = \psi_0(x_1)\bar{\psi}_0(x_2)S^{nz}(x_3, x_4)\dots S^{nz}(x_{2m-1}, x_{2m})det'(\rho). \end{aligned} \quad (11)$$

3 Multi-Fermion Contribution to the Observable Values

Since boson part of the amplitude of the process (3) is well known in Gauss approximation let us concentrate on the fermion contribution only and consider for shortening that there are not gauge and Higgs bosons in final state

$$f + f \rightarrow (2m - 2)f. \quad (12)$$

Generalization into arbitrary number of boson is trivial. For calculation of the amplitude we can apply standard Lehman-Symanzik-Zimmermann technique:

$$\begin{aligned} A(k_1, p_1, \dots, k_m, p_m) &= \int_0^\infty \frac{d\rho}{\rho^5} \mu(\rho) e^{-\frac{8\pi^2}{g^2} - \pi^2 \rho^2 v^2} \tilde{\psi}_0(k_1) \tilde{\bar{\psi}}_0(p_1) \times \\ &\times k_{1\eta} \sigma_\eta p_{1\eta'} \bar{\sigma}_{\eta'} det'(\rho) \prod_{j=2}^m \tilde{S}^{nz}(p_j, k_j) k_{j\mu} \sigma_\mu p_{j\nu} \bar{\sigma}_\nu \delta^{(4)}\left(p_1 + k_1 - \sum_{n=2}^m (p_n + k_n)\right), \end{aligned} \quad (13)$$

where k_1 and p_1 denotes 4-momenta of initial particles, $k_2, p_2, \dots, k_m, p_m$ are 4-momenta of final fermions, tildes mean Fourier transformations of corresponding functions, δ -function appears due to the integration over instanton size z_μ . It is supposed that all momenta satisfy the mass-shell equation.

The contribution of the initial bosons and zero modes are well known. Moreover, as is shown below, the leading fermion contribution into cross-section comes from non-zero modes. Therefore we restrict ourselves to non-zero modes contribution into the amplitude (13). First of all Fourier transformation of non-zero modes propagator (9) have to be done

$$\tilde{S}^{nz}(p, k) \equiv \int dx dy e^{-ipx - ipy} S^{nz}(x, y) = -2\pi^2 \rho^2 (p_\mu \bar{\sigma}_\mu + k_\nu \sigma_\nu) \frac{p_\lambda \sigma_\lambda k_{\lambda'} \bar{\sigma}_{\lambda'}}{p^2 k^2 (pk)}, \quad (14)$$

where we expand the expression in powers of ρ^2 and neglect terms of order $O(\rho^4)$ [11]. Amputation of external legs and going to mass-shell gives

$$\lim_{p^2, k^2 \rightarrow 0} \hat{S}^{nz}(p, k) k_\eta \sigma_\eta p_{\eta'} \bar{\sigma}_{\eta'} = -2\pi^2 \rho^2 \frac{p_\mu \bar{\sigma}_\mu + k_\nu \sigma_\nu}{(pk)}. \quad (15)$$

Substituting (15) into (13) and performing integration over ρ we obtain

$$\begin{aligned} A(k_1, p_1, \dots, k_m, p_m) &= C(p_1, k_1) (m-1)! \times \\ &\times \prod_{j=2}^m \left(\frac{-2}{v^2} \right) \frac{p_{j\mu} \bar{\sigma}_\mu + k_{j\nu} \sigma_\nu}{(p_j k_j)} \delta^{(4)}\left(p_1 + k_1 - \sum_{n=2}^m (p_n + k_n)\right), \end{aligned} \quad (16)$$

where zero modes are contained in factor C . Squaring and integrating over 3-momenta of final particles we obtain the probability of m fermion pairs production in instanton field

$$\begin{aligned}
 P_m &= \int_0^{\sqrt{s}} \frac{d\vec{p}_1}{(2\pi)^3} \frac{d\vec{k}_1}{(2\pi)^3} |C(p_1, k_1)|^2 (m-1)! \prod_{j=2}^m \frac{4}{v^4} \int_0^{\sqrt{s}} \frac{d\vec{p}_j}{(2\pi)^3} \frac{d\vec{k}_j}{(2\pi)^3} (p_j k_j)^{-1} \times \\
 &\times \delta^{(4)}\left(p_1 + k_1 - \sum_{n=2}^m (p_n + k_n)\right) = \frac{N}{(m-1)!} \left(\frac{4}{v^4} 4\pi \left(\frac{1 - \ln 2}{3(2\pi)^6} \right) \frac{\sqrt{s}^6}{m_0^2} \right)^{m-1} = \\
 &= \frac{N}{(m-1)!} \left(\frac{\sqrt{s}^6}{m_0^2 m_w^2 E_{sp}^2} \frac{6(1 - \ln 2)}{\pi^3} \right)^{m-1}, \tag{17}
 \end{aligned}$$

where small mass parameter m_0 is introduced for regularization of the collinear singularity, condition $m_0 \ll \sqrt{s}$ is used, factor N guarantees the probability normalization $\sum_{m=1}^{\infty} P_m = 1$:

$$N = \exp \left[-\frac{\sqrt{s}^6}{m_0^2 m_w^2 E_{sp}^2} \frac{6(1 - \ln 2)}{\pi^3} \right]. \tag{18}$$

The contribution of fermions into total instanton-induced cross-section with exponential accuracy is

$$\sigma_f^{inst} \sim \exp \left[\frac{n_f \sqrt{s}^6}{m_0^2 m_w^2 E_{sp}^2} \frac{6(1 - \ln 2)}{\pi^3} \right], \tag{19}$$

where we took into account arbitrary number of fermions sorts. Fig.1 shows the dependence of $\ln \sigma_f^{inst}$ on energy.




Fig.1. Logarithm of the fermion part of the total instanton-induced cross-section in dependence on energy. We took for definiteness $m_w = 80 \text{ GeV}$, $m_0 = 0.5 \text{ MeV}$, $n_f = 12$.

4 Conclusion

We obtain Poisson distribution on number of fermion pairs (17) produced in the instanton processes. Average number of such pairs reads

$$\langle m \rangle = \frac{n_f \sqrt{s}^6}{m_0^2 m_w^2 E_{sp}^2} \frac{6(1 - \ln 2)}{\pi^3}. \quad (20)$$

Fermion part of instanton cross section (19) strongly (exponentially) increases with energy. The contribution of zero modes is included in pre-exponent factor. Non-zero fermion modes are accumulated in exponent index and play essential role. At low energies contribution (19) is not important. At high energies $\sqrt{s} \sim \sqrt[3]{m_w m_0 E_{sp}} \sim O(10) \text{ GeV}$ exponent index is equal approximately to unit (see Fig.1). Of course, at such energies fermion contribution is not essential due to suppressing 't Hooft factor $\exp[-\frac{16\pi^2}{g^2}] \approx 10^{-169}$. But at energies $\sqrt{s} \sim E_{sp}$ one can suppose that 't Hooft suppression disappears and then strongly rising fermion exponent have to be taken into account.

Fermion contribution into instanton-induced cross-section calculated by means of coherent-state formalism by O.Espinosa [11] also exponentially increase with energy $\sigma_f^{inst} \sim \exp[\sqrt{s}^{4/3}]$. In comparison with the result of Ref. [11] straightforward method gives $(\sqrt{s})^6$ in exponent index. The difference can be caused the approximations inherent to coherent-state method for fermionic degrees of freedom.

References

- [1] G.'t Hooft. Phys.Rev.Lett. **37** 8 (1976), Phys.Rev. **D14** 3432 (1976).
- [2] A.Sakharov. JETP Lett. **5** 1 (1967).
- [3] A.Ringwald. Nucl.Phys. **B330** 1 (1990).
- [4] M.Porrati. Nucl.Phys. **B347** 371 (1990).
- [5] V.Zakharov. Nucl.Phys. **B371** 637 (1992).
- [6] M.Mattis. Phys.Rep. **214** 159 (1992).
- [7] N.Manton. Phys.Rev. **D28** 2019 (1983); F.Klinhamer, N.Manton. Phys.Rev. **D30** 2212 (1984).
- [8] V.Khoze, A.Ringwald. Nucl.Phys. **B355** 351 (1991).
- [9] D.Diakonov, V.Petrov. Proc. of the XXVI LNPI Winter School, Leningrad 1991, 8 (1991), in Russian.
- [10] S.Khlebnikov, V.Rubakov, P.Tinyakov. Nucl.Phys. **B350** 441 (1991).
- [11] O.Espinosa. Nucl.Phys. **B375** 263 (1992).
- [12] A.Vainstein et al. Usp.Fiz.Nauk **13** 553 (1982), in Russian.
- [13] A.Belavin, A.Polyakov, A.Schwarz and Yu.Tyupkin. Phys.Lett. **B59** 85 (1975).
- [14] L.Brown et al. Phys.Rev. **D17** 1583 (1978).
- [15] T.Schaefer, E.Shuryak. Rev.Mod.Phys. **70** 323 (1998).